

REHAB QUIZ

1. **All are electrophysiological diagnostic criteria of carpal tunnel syndrome except**
 - A) Distal median motor latency > 4.4 Ms
 - B) Difference between distal motor latency of median and ulnar > 0.4 Ms
 - C) Difference between distal sensory latency of median and ulnar > 0.2 Ms
 - D) Inching technique: latency jump > 0.2 Ms/cm

2. **Commonest inspiratory muscle training method prescribed in patients with COPD**
 - A) Threshold loading technique
 - B) Flow resistive technique
 - C) Purse lip breathing technique
 - D) Diaphragmatic breathing technique

3. **Brandt-Daroff exercise is used for**
 - A) In hospital treatment for cervical vertigo
 - B) Home exercise for cervical vertigo
 - C) In hospital treatment for BPPV
 - D) Home exercise for BPPV

4. **Which artery is involved in a stroke patient presented with contralateral hemianaesthesia with visual special deficit and aprosody without affect disorder**
 - A) Upper division of MCA – dominant hemisphere
 - B) Upper division of MCA – non-dominant hemisphere
 - C) Lower division of MCA – dominant hemisphere
 - D) Lower division of MCA – non-dominant hemisphere

5. **Which lobe of lung is drained in supine position with foot end of bed elevation 18 – 20 inches?**
 - A) Anterior basal segment of right lower lobe
 - B) Lateral basal segment of right lower lobe
 - C) Medial basal segment of right lower lobe
 - D) Lateral basal segment of left lower lobe

6. **Both chin tuck and supraglottic swallow is helpful in**
 - A) Delayed swallowing reflex and decreased unilateral pharyngeal paralysis
 - B) Delayed swallowing reflex and decreased bilateral pharyngeal paralysis
 - C) Delayed swallowing reflex and decreased laryngeal closure
 - D) Delayed swallowing reflex and decreased opening of cricopharyngeal region

PG Forum

7. All are energy storing prosthetic foot except
- A) SAFE II foot
 - B) Seattle foot
 - C) Carbon copy II foot
 - D) Flex foot
8. Calf hypertrophy is not seen in
- A) Duchene muscular dystrophy
 - B) Becker muscular dystrophy
 - C) Limb girdle muscular dystrophy
 - D) Emery- Dreifuss muscular dystrophy
9. Modified Amor criteria is diagnostic of
- A) Bechet's disease
 - B) Undifferentiated spondylo-arthropathy
 - C) Juvenile onset spondylo-arthropathy
 - D) Electrophysiological criteria of GBS
10. All are done in speech therapy of aphasia except
- A) Melody intonation technique
 - B) Retraining of articular pattern
 - C) Conversational coaching
 - D) Oral reading.

ANSWERS

June issue:

1-C; 2-D; 3-D; 4-C; 5-A; 6-D; 7-B; 8-A; 9-C; 10-C

Right responder (10 out of 10): **Dr. Harleen Uppal**, Junior Resident, Dept. of PMR, AIIMS, New Delhi

Next right responder: **Dr. Prajna Ranjani M**, Junior Resident, AIIMS, New Delhi