

## Rehab Quiz

1. Which is TRUE? Coordination of micturition involves
  - a. The sacral micturition centre in which the afferent parasympathetic impulses provide feedback on bladder fullness
  - b. The brainstem that coordinates contraction of the bladder with relaxation of the sphincter
  - c. The cerebral cortex that prevents incontinence of urine during forceful activities such as coughing
  - d. The pontine centre that exerts the final control to initiate or delay voiding
2. Regarding CRP, which of the following statements is not correct?
  - a. Increased in any inflammation due to trauma, infection, malignancies, etc
  - b. In SLE elevated CRP suggests infection or serositis
  - c. It follows rise and fall in ESR
  - d. It is markedly elevated in Still's disease
3. Normal cross section of median nerve in CTS by USG at wrist crease is
  - a. 9–11 mm
  - b. 11–13 mm
  - c. 13–15 mm
  - d. 15–19 mm
4. What is the Stimson technique?
  - a. A provocative maneuver to test for glenohumeral instability
  - b. Gravity-assisted technique to reduce an anterior shoulder dislocation
  - c. A two-person technique to reduce an anterior shoulder dislocation, using a sheet under the axilla by one person and manual traction by the other person
  - d. A test for inferior shoulder laxity
5. Electrodiagnostic findings common in myopathies include:
  - a. Low amplitude sensory nerve action potentials (SNAP)
  - b. Long duration motor units
  - c. Abnormal spontaneous potentials (denervation) on needle EMG
  - d. Increased firing frequency of motor units
6. A patient with a brainstem lesion is likely to experience all of the following except:
  - a. Poor anterior-posterior transport of a food/liquid bolus
  - b. Decreased masseter tension
  - c. Wet hoarseness
  - d. Rigidity
7. Which of the following is considered a poor prognostic factor in patients with amyotrophic lateral sclerosis (ALS)?
  - a. Predominance of upper motor neuron (UMN) findings at diagnosis
  - b. Long period from symptom onset to diagnosis
  - c. Younger age of onset
  - d. Pulmonary dysfunction early in the clinical course
8. The first line of treatment for bothersome phantom limb sensation, phantom limb pain and residual limb pain is:
  - a. Desensitization techniques
  - b. Pharmacologic techniques
  - c. Biofeedback techniques
  - d. Use of transcutaneous electrical nerve stimulation (TENS) unit
9. Which statement is TRUE regarding urinary tract infection (UTI) in individuals with spinal cord injury?
  - a. The NIDRR consensus definition of UTI requires the presence of significant bacteriuria, pyuria or signs and symptoms
  - b. The presenting symptoms can be that of autonomic dysreflexia
  - c. Asymptomatic bacteriuria in patients performing intermittent catheterization should be treated
  - d. Urine for urinalysis and culture should be collected from the collecting device just prior to changing the catheter
10. A patient presents for electrodiagnostic testing for left-sided low back pain. The study is normal except for a prolonged left H-reflex. What is your diagnosis?
  - a. Left S1 radiculopathy
  - b. Polyneuropathy
  - c. Lumbar plexopathy
  - d. None of the above